

Lattice Laws Forcing Distributivity Under Unique Complementation

*R. Padmanabhan**

Department of Mathematics
University of Manitoba
Winnipeg, Manitoba R3T 2N2
Canada

W. McCune†

Mathematics and Computer Science Division
Argonne National Laboratory
Argonne, Illinois 60439-4844
U.S.A.

Robert Veroff‡

Department of Computer Science
University of New Mexico
Albuquerque, New Mexico 87131
U.S.A.

August 8, 2005

Abstract

In this paper we give several new lattice identities valid in non-modular lattices such that a uniquely complemented lattice satisfying any of these identities is necessarily Boolean. Since some of these identities are consequences of modularity as well, these results generalize the classical result of Birkhoff and von Neumann that every uniquely complemented modular lattice is Boolean. In particular, every uniquely complemented lattice in $M \vee N_5$, the least non-modular variety, is Boolean.

*Supported by an operating grant from NSERC of Canada (OGP8215).

†Supported by the Mathematical, Information, and Computational Sciences Division sub-program of the Office of Advanced Scientific Computing Research, U.S. Department of Energy, under Contract W-31-109-Eng-38.

‡Supported in part by . . .

1 Introduction

In 1904 Huntington [4] conjectured that every uniquely complemented lattice must be distributive (and hence a Boolean algebra). In 1945, R. P. Dilworth shattered this conjecture by proving [2] that every lattice can be embedded in a uniquely complemented lattice. For a much powerful version of the same result, see Adams and Sichler [1].

In spite of these deep results, still it is hard to find "nice" examples of uniquely complemented lattices that are not Boolean. This is because uniquely complemented lattices having a little extra structure most often turn out to be distributive. This seems to be the essence of Huntington's conjecture. For example, we have the theorem of Garrett Birkhoff and von Neumann that every uniquely complemented modular lattice is Boolean. Following [8], we call a lattice property P a *Huntington property* if every uniquely complemented P -lattice is distributive. Similarly, a lattice variety K is said to be a *Huntington variety* if every uniquely complemented lattice in K is Boolean. In this terminology, the modular lattices are the largest known Huntington variety. A monograph by Salii [11] gives a comprehensive survey of known Huntington properties. Among these, modularity is the only known condition which is a lattice identity. In this paper, we give a number of new non-modular Huntington varieties and any of them could be construed as a generalization of von Neumann-Birkhoff theorem.

The automated theorem provers Otter [5] and Prover9 [7], and the program Mace4 [6], which searches for finite algebras, were used in this work. Several automated proofs are given in an appendix to this paper. The Web page associated with this paper [10] contains additional Huntington identities and additional automated proofs supporting this work.

2 A Non-modular Huntington Variety

Here we give a lattice identity that defines a non-modular Huntington variety. Several others are given in the following sections and in the supporting Web page [10].

Theorem 1. *The variety of lattices defined by*

$$(x \wedge (y \vee (x \wedge z))) \vee (x \wedge (z \vee (x \wedge y))) = x \wedge (z \vee y) \quad (\text{H69})$$

is a non-modular Huntington variety.

Proof. We show that the condition $a \wedge b' = 0$ forces the inequality $a \leq b$ and hence by a well-known theorem of O. Frink (e.g. see [9]), the lattice will necessarily be Boolean. Indeed, let $a \wedge b' = 0$ for some two elements a, b in a uniquely complemented lattice satisfying the identity

$$(x \wedge (y \vee (x \wedge z))) \vee (x \wedge (z \vee (x \wedge y))) = x \wedge (z \vee y).$$

Put $z = x'$ in the above to get

$$(x \wedge y) \vee (x \wedge (x' \vee (x \wedge y))) = x \wedge (x' \vee y).$$

Now let $x = b'$, $y = a$. We have

$$(b' \wedge a) \vee (b' \wedge (b \vee (b' \wedge a))) = b' \wedge (b \vee a).$$

So if we assume that $a \wedge b' = 0$, then we get $b' \wedge (b \vee a) = 0$. Also, $b' \vee (b \vee a) = (b' \vee b) \vee a = 1 \vee a = 1$. Thus both b and $b \vee a$ are complements of the element b' . Since the lattice is uniquely complemented, we get the desired conclusion $b \vee a = b$. In other words, we have proved that the given lattice satisfies the bi-implication $a \leq b$ if and only if $a \wedge b' = 0$ and hence, by Frink's theorem, the lattice is distributive. $\square$

3 Huntington Implications

Here we show Huntington properties that are implications. These can be used, among other purposes, to show that lattice identities are Huntington.

Theorem 2. (See [8].) *A uniquely complemented lattice satisfying any one of the following three implications (or their duals) is distributive.*

$$x \vee y = x \vee z \Rightarrow x \vee y = x \vee (y \wedge z) \quad (\text{SD-}\vee)$$

$$x \vee y = x \vee z \Rightarrow (x \wedge y) \vee (x \wedge z) = x \wedge (y \vee z) \quad (\text{CD-}\vee)$$

$$x \vee y = x \vee z \Rightarrow x \wedge ((x \wedge y) \vee z) = (x \wedge y) \vee (x \wedge z) \quad (\text{CM-}\vee)$$

A proof of (CD- $\vee$) is given in the appendix. Proofs of the other two cases are given on the supporting Web page [10].

Corollary 1. *A uniquely complemented lattice satisfying the identity*

$$x \wedge ((y \wedge (x \vee z)) \vee (z \wedge (x \vee y))) = (x \wedge y) \vee (x \wedge z) \quad (\text{H82})$$

is Huntington.

Proof. It is easy to see that (H82) implies the lattice implication (CD- $\vee$). Indeed, if

$$x \vee y = x \vee z,$$

then

$$\begin{aligned} (x \wedge y) \vee (x \wedge z) &= x \wedge ((y \wedge (x \vee z)) \vee (z \wedge (x \vee y))) && \text{by (H82)} \\ &= x \wedge ((y \wedge (x \vee y)) \vee (z \wedge (x \vee z))) && \text{by hypothesis} \\ &= x \wedge (y \vee z) \end{aligned}$$

$\square$

As the reader can see, the identity (H82) is designed to show that there are lattice identities which formally imply such implications. Using powerful concepts like the bounded homomorphisms of Ralph McKenzie, one could show

that there many lattice identities like (H82) which formally imply (SD- $\vee$), (SD- $\wedge$), (CD- $\vee$), etc. In fact, every finite lattice satisfying (SD- $\vee$) or (SD- $\wedge$) will satisfy a lattice identity which formally implies the respective implication and all these identities are examples of non-modular Huntington identities (for more details, please see [8]).

Table 1 lists several Huntington identities justified by the preceding Huntington implications. Proofs can be found on the supporting Web page [10]. None of the identities are equivalent (given lattice theory).

Name	Identity	Reason
H18	$(x \wedge y) \vee (x \wedge z) = x \wedge ((x \wedge y) \vee ((x \wedge z) \vee (y \wedge (x \vee z))))$	CM- $\vee$
H50	$x \wedge (y \vee (z \wedge (x \vee u))) = x \wedge (y \vee (z \wedge (x \vee (z \wedge (y \vee u)))))$	SD- $\vee$
H51	$x \wedge (y \vee (z \wedge (x \vee u))) = x \wedge (y \vee ((x \wedge z) \vee (z \wedge u)))$	SD- $\vee$
H64	$x \wedge (y \vee z) = x \wedge (y \vee (x \wedge (z \vee (x \wedge (y \vee (x \wedge z)))))$	SD- $\wedge$
H68	$x \wedge (y \vee z) = x \wedge (y \vee (x \wedge (z \vee (x \wedge y))))$	SD- $\wedge$
H69	$x \wedge (y \vee z) = (x \wedge (z \vee (x \wedge y))) \vee (x \wedge (y \vee (x \wedge z)))$	SD- $\wedge$
H76	$x \wedge (y \vee (z \wedge (y \vee u))) = x \wedge (y \vee (z \wedge (u \vee (x \wedge y))))$	SD- $\vee$, SD- $\wedge$
H79	$x \wedge (y \vee (z \wedge (x \vee u))) = x \wedge ((x \wedge (y \vee (x \wedge z))) \vee (z \wedge u))$	SD- $\vee$, SD- $\wedge$
H80	$(x \wedge y) \vee (x \wedge z) = x \wedge ((x \wedge y) \vee (z \wedge (x \vee (y \wedge (x \vee z)))))$	CM- $\vee$
H82	$(x \wedge y) \vee (x \wedge z) = x \wedge ((y \wedge (x \vee z)) \vee (z \wedge (x \vee y)))$	CD- $\vee$, CM- $\vee$

Table 1: Huntington Identities Justified by Huntington Implications

4 More Huntington Identities

This section contains several non-modular Huntington identities that do not satisfy the Huntington implications (SD- $\vee$), (CD- $\vee$), (CM- $\vee$), or their duals.

Theorem 3. *The variety of lattices defined by*

$$x \wedge (y \vee z) = x \wedge (y \vee ((x \vee y) \wedge (z \vee (x \wedge y)))) \quad (\text{H58})$$

is a non-modular Huntington variety.

Proof. (The automatic proof from which this proof was derived is given in the Appendix.) We show that any uniquely complemented lattice satisfying (H58) also satisfies the property $a \leq b \Rightarrow b' \leq a'$. Assume $a \leq b$ and therefore $a \wedge b' = 0$. In (H58) set $x = a, y = b, z = (a \wedge b')'$ and then simplify, giving

$$a \wedge (b' \vee (a \vee b')') = 0.$$

Then unique complementation gives $b' \vee (a \vee b')' = a'$ and therefore $b' \leq a'$. $\square$

Additional Huntington identities not satisfying the Huntington implications are shown in the following list. Automated proofs are given on the supporting

RP: what is the name of this property? Should we justify using it?

Web page [10].

$$x \wedge (y \vee (z \wedge (x \vee u))) = x \wedge (y \vee (z \wedge (x \vee (z \wedge u)))) \quad (\text{H1})$$

$$x \wedge (y \vee (x \wedge z)) = x \wedge (y \vee (z \wedge ((x \wedge (y \vee z)) \vee (y \wedge z)))) \quad (\text{H2})$$

$$x \wedge (y \vee (x \wedge z)) = x \wedge (y \vee (z \wedge (y \vee (x \wedge (z \vee (x \wedge y)))))) \quad (\text{H3})$$

$$x \vee (y \wedge (x \vee z)) = x \vee (y \wedge (z \vee (x \wedge (z \vee y)))) \quad (\text{H55})$$

$$x \wedge (y \vee z) = x \wedge (y \vee ((x \vee y) \wedge (z \vee (x \wedge y)))) \quad (\text{H58})$$

References

- [1] M. E. Adams and J. Sichler. Lattice with unique complementation. *Pac. J. Math.*, 92:1–13, 1981.
- [2] R. P. Dilworth. Lattices with unique complements. *Trans. AMS*, 57:123–154, 1945.
- [3] G. Grätzer. *General Lattice Theory*. Birkhauser Verlag, 1975.
- [4] E. V. Huntington. Sets of independent postulates for the algebra of logic. *Trans. AMS*, 5:288–309, 1904.
- [5] W. McCune. Otter 3.0 Reference Manual and Guide. Tech. Report ANL-94/6, Argonne National Laboratory, Argonne, IL, 1994. Also see <http://www.mcs.anl.gov/AR/otter/>.
- [6] W. McCune. Mace4 Reference Manual and Guide. Tech. Memo ANL/MCS-TM-264, Mathematics and Computer Science Division, Argonne National Laboratory, Argonne, IL, August 2003.
- [7] W. McCune. Prover9. <http://www.mcs.anl.gov/~mccune/prover9/>, 2005.
- [8] R. Padmanabhan. Huntington varieties of lattices. To Appear.
- [9] R. Padmanabhan. A first-order proof of a theorem of Frink. *Algebra Universalis*, 13(3):397–400, 1981.
- [10] R. Padmanabhan, W. McCune, and R. Veroff. Lattice laws forcing distributivity under unique complementation: Web support. <http://www.mcs.anl.gov/~mccune/papers/uc-lattice/>, 2005.
- [11] V. N. Salii. Lattices with unique complements. *Trans. AMS*, 69, 1988.

Appendix

Covers of N_5

We should include some introduction to these lattices.




Figure 1: All Covers of the Least Non-Modular Lattice N_5




Figure 2: Lattices M_3 , N_5 , $NM08$

(H1)	1	2				7			10	11		13	14	15		N_5
(H2)	1		4		6	7	8	9	10	11	12	13	14	15	M_3	N_5 NM08
(H3)	1		4		6	7	8	9	10	11	12	13	14	15	M_3	N_5 NM08
(H18)	1		4		6	7		9	10	11		13		15	M_3	N_5 NM08
(H50)	1				6	7		9	10	11		13	14	15		N_5
(H51)	1					7			10	11		13	14	15		N_5
(H55)		2			5	6	7	8	9	10		12	13	14	15	M_3 N_5 NM08
(H58)		2			5	6	7	8	9	10		12	13	14	15	M_3 N_5 NM08
(H64)		2				6	7	8	9	10	11	12	13	14	15	N_5
(H68)		2				6	7	8	9	10		12	13	14	15	N_5
(H69)		2				6	7	8	9	10		12	13	14	15	N_5
(H76)						6	7	8	9	10		13	14	15		N_5
(H79)							7			10		13	14	15		N_5
(H80)	1		3	4		6	7	8	9	10	11	13		15	M_3	N_5 NM08
(H82)	1			4		6	7		9	10	11	13		15		N_5

Table 2: Lattices ($L_1 - L_{15}$, M_3 , N_5 , NM08) for which the Identities Hold**Proof of Theorem 2, Part CD- $\vee$**

This proof was produced by the program Prover9 [7]. The input and output files can be found on the supporting Web page [10].

13	$x \vee y = y \vee x$	[input]
14	$x \wedge y = y \wedge x$	[input]
15	$(x \vee y) \vee z = x \vee (y \vee z)$	[input]
16	$(x \wedge y) \wedge z = x \wedge (y \wedge z)$	[input]
17	$x \wedge (x \vee y) = x$	[input]
18	$x \vee (x \wedge y) = x$	[input]
19	$x \vee x' = 1$	[input]
20	$x \wedge x' = 0$	[input]
21	$x \vee y \neq 1 \mid x \wedge y \neq 0 \mid x' = y$	[input]
22	$A \wedge B = A$	[input]
23	$A' \vee B' \neq A'$	[input]
24	$x \vee y \neq x \vee z \mid x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$	[input]
26	$x \wedge (y \wedge z) = y \wedge (x \wedge z)$	[14 $\rightarrow$ 16; 16]
32	$x \vee ((x \wedge y) \vee z) = x \vee z$	[18 $\rightarrow$ 15]
37	$x \vee (x' \vee y) = 1 \vee y$	[19 $\rightarrow$ 15]
39	$x \wedge 1 = x$	[19 $\rightarrow$ 17]
42	$x \vee 0 = x$	[20 $\rightarrow$ 18]
43	$A \wedge (B \wedge x) = A \wedge x$	[22 $\rightarrow$ 16]
51	$x \vee y \neq 1 \mid x \wedge (y \vee x') = 0 \vee (x \wedge y)$	[19 $\rightarrow$ 24; 20 13]
60	$1 \wedge x = x$	[39 $\rightarrow$ 14]
66	$0 \vee x = x$	[42 $\rightarrow$ 13]
69	$x \vee y \neq 1 \mid x \wedge (y \vee x') = x \wedge y$	[51; 66]
72	$1 \vee x = 1$	[60 $\rightarrow$ 17]
73	$x \vee (x' \vee y) = 1$	[37; 72]

75	$0 \wedge x = 0$	[66 $\rightarrow$ 17]
79	$x \wedge (y \wedge x') = y \wedge 0$	[20 $\rightarrow$ 26]
81	$x \vee 1 = 1$	[72 $\rightarrow$ 13]
83	$x \wedge 0 = 0$	[75 $\rightarrow$ 14]
84	$x \wedge (y \wedge x') = 0$	[79; 83]
103	$A \wedge (A' \vee B') \neq 0$	[21 a 73 a c 23 a]
170	$x \vee (x \wedge y)' = 1$	[19 $\rightarrow$ 32; 81]
185	$x \vee (y \wedge x)' = 1$	[14 $\rightarrow$ 170]
194	$B \vee A' = 1$	[22 $\rightarrow$ 185]
891	$B \wedge (A' \vee B') = B \wedge A'$	[69 a 194 a]
5852	$A \wedge (A' \vee B') = 0$	[891 $\rightarrow$ 43; 84]
5853	$\square$	[5852 a 103 a]

Proof of Theorem 3

This proof was produced by the program Prover9 [7]. The input and output files can be found on the supporting Web page [10].

29	$x \vee y = y \vee x$	[input]
37	$x \vee (y \wedge x) = x$	[input]
38	$x \vee (x \vee y) = x \vee y$	[input]
40	$x \wedge x' = 0$	[input]
47	$x \vee 0 = x$	[input]
48	$0 \vee x = x$	[input]
49	$x \vee y \neq 1 \mid x \wedge y \neq 0 \mid x' = y$	[input]
50	$x \wedge (x \vee y)' = 0$	[input]
51	$x \vee (y \vee (x \vee y)') = 1$	[input]
52	$x \wedge (y \vee ((x \vee y) \wedge (z \vee (x \wedge y)))) = x \wedge (y \vee z)$	[input]
53	$A \wedge B = A$	[input]
54	$A' \vee B' \neq A'$	[input]
103	$A \vee B = B$	[53 $\rightarrow$ 37; 29]
107	$A \wedge B' = 0$	[103 $\rightarrow$ 50]
109	$A \wedge (B' \vee ((A \vee B') \wedge x)) = A \wedge (B' \vee x)$	[107 $\rightarrow$ 52; 47]
2392	$A \wedge (B' \vee (A \vee B')') = 0$	[40 $\rightarrow$ 109; 29 48 107]
2439	$B' \vee (A \vee B')' = A'$	[49 a 51 a b 2392 a]
2481	$A' \vee B' = A'$	[2439 $\rightarrow$ 38; 29 2439]
2482	$\square$	[2481 a 54 a]