

## Annihilator Method

### CS 461, Lecture 4

Jared Saia  
University of New Mexico

- Write down the annihilator for the recurrence
- Factor the annihilator
- Look up the factored annihilator in the “Lookup Table” to get general solution
- Solve for constants of the general solution by using initial conditions

2

### Today's Outline

### Lookup Table

- Annihilators for recurrences with non-homogeneous terms
- Transformations

$$(\mathbf{L} - a_0)^{b_0}(\mathbf{L} - a_1)^{b_1} \dots (\mathbf{L} - a_k)^{b_k}$$

annihilates only sequences of the form:

$$\langle p_1(n)a_0^n + p_2(n)a_1^n + \dots p_k(n)a_k^n \rangle$$

where  $p_i(n)$  is a polynomial of degree  $b_i - 1$  (and  $a_i \neq a_j$ , when  $i \neq j$ )

1

3

## Examples

- Q: What does  $(L - 3)(L - 2)(L - 1)$  annihilate?
- A:  $c_0 1^n + c_1 2^n + c_2 3^n$
- Q: What does  $(L - 3)^2(L - 2)(L - 1)$  annihilate?
- A:  $c_0 1^n + c_1 2^n + (c_2 n + c_3) 3^n$
- Q: What does  $(L - 1)^4$  annihilate?
- A:  $(c_0 n^3 + c_1 n^2 + c_2 n + c_3) 1^n$
- Q: What does  $(L - 1)^3(L - 2)^2$  annihilate?
- A:  $(c_0 n^2 + c_1 n + c_2) 1^n + (c_3 n + c_4) 2^n$

4

## Example

Consider the recurrence  $T(n) = 7T(n - 1) - 16T(n - 2) + 12T(n - 3)$ ,  $T(0) = 1$ ,  $T(1) = 5$ ,  $T(2) = 17$

- **Write down the annihilator:** From the definition of the sequence, we can see that  $L^3 T - 7L^2 T + 16L T - 12T = 0$ , so the annihilator is  $L^3 - 7L^2 + 16L - 12$
- **Factor the annihilator:** We can factor by hand or using a computer program to get  $L^3 - 7L^2 + 16L - 12 = (L - 2)^2(L - 3)$
- **Look up to get general solution:** The annihilator  $(L - 2)^2(L - 3)$  annihilates sequences of the form  $\langle (c_0 n + c_1) 2^n + c_2 3^n \rangle$
- **Solve for constants:**  $T(0) = 1 = c_1 + c_2$ ,  $T(1) = 5 = 2c_0 + 2c_1 + 3c_2$ ,  $T(2) = 17 = 8c_0 + 4c_1 + 9c_2$ . We've got three equations and three unknowns. Solving by hand, we get that  $c_0 = 1, c_1 = 0, c_2 = 1$ . **Thus:**  $T(n) = n2^n + 3^n$

5

## Example (II)

Consider the recurrence  $T(n) = 2T(n - 1) - T(n - 2)$ ,  $T(0) = 0$ ,  $T(1) = 1$

- **Write down the annihilator:** From the definition of the sequence, we can see that  $L^2 T - 2L T + T = 0$ , so the annihilator is  $L^2 - 2L + 1$
- **Factor the annihilator:** We can factor by hand or using the quadratic formula to get  $L^2 - 2L + 1 = (L - 1)^2$
- **Look up to get general solution:** The annihilator  $(L - 1)^2$  annihilates sequences of the form  $(c_0 n + c_1) 1^n$
- **Solve for constants:**  $T(0) = 0 = c_1$ ,  $T(1) = 1 = c_0 + c_1$ . We've got two equations and two unknowns. Solving by hand, we get that  $c_0 = 0, c_1 = 1$ . **Thus:**  $T(n) = n$

6

## At Home Exercise

Consider the recurrence  $T(n) = 6T(n - 1) - 9T(n - 2)$ ,  $T(0) = 1$ ,  $T(1) = 6$

- Q1: What is the annihilator of this sequence?
- Q2: What is the factored version of the annihilator?
- Q3: What is the general solution for the recurrence?
- Q4: What are the constants in this general solution?

(Note: You can check that your general solution works for  $T(2)$ )

7

## Non-homogeneous terms

- Consider a recurrence of the form  $T(n) = T(n-1) + T(n-2) + k$  where  $k$  is some constant
- The terms in the equation involving  $T$  (i.e.  $T(n-1)$  and  $T(n-2)$ ) are called the *homogeneous* terms
- The other terms (i.e.  $k$ ) are called the *non-homogeneous* terms

8

## Example

- In a *height-balanced tree*, the height of two subtrees of any node differ by at most one
- Let  $T(n)$  be the smallest number of nodes needed to obtain a height balanced binary tree of height  $n$
- Q: What is a recurrence for  $T(n)$ ?
- A: Divide this into smaller subproblems
  - To get a height-balanced tree of height  $n$  with the smallest number of nodes, need one subtree of height  $n-1$ , and one of height  $n-2$ , plus a root node
  - Thus  $T(n) = T(n-1) + T(n-2) + 1$

9

## Example

- Let's solve this recurrence:  $T(n) = T(n-1) + T(n-2) + 1$  (Let  $T_n = T(n)$ , and  $T = \langle T_n \rangle$ )
- We know that  $(\mathbf{L}^2 - \mathbf{L} - 1)$  annihilates the homogeneous terms
- Let's apply it to the entire equation:

$$\begin{aligned}(\mathbf{L}^2 - \mathbf{L} - 1)\langle T_n \rangle &= \mathbf{L}^2\langle T_n \rangle - \mathbf{L}\langle T_n \rangle - 1\langle T_n \rangle \\ &= \langle T_{n+2} \rangle - \langle T_{n+1} \rangle - \langle T_n \rangle \\ &= \langle T_{n+2} - T_{n+1} - T_n \rangle \\ &= \langle 1, 1, 1, \dots \rangle\end{aligned}$$

10

## Example

- This is close to what we want but we still need to annihilate  $\langle 1, 1, 1, \dots \rangle$
- It's easy to see that  $\mathbf{L} - 1$  annihilates  $\langle 1, 1, 1, \dots \rangle$
- Thus  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 1)$  annihilates  $T(n) = T(n-1) + T(n-2) + 1$
- When we factor, we get  $(\mathbf{L} - \phi)(\mathbf{L} - \hat{\phi})(\mathbf{L} - 1)$ , where  $\phi = \frac{1+\sqrt{5}}{2}$  and  $\hat{\phi} = \frac{1-\sqrt{5}}{2}$ .

11

## Lookup

- Looking up  $(\mathbf{L} - \phi)(\mathbf{L} - \hat{\phi})(\mathbf{L} - 1)$  in the table
- We get  $T(n) = c_1\phi^n + c_2\hat{\phi}^n + c_31^n$
- If we plug in the appropriate initial conditions, we can solve for these three constants
- We'll need to get equations for  $T(2)$  in addition to  $T(0)$  and  $T(1)$

12

## General Rule

To find the annihilator for recurrences with non-homogeneous terms, do the following:

- Find the annihilator  $a_1$  for the homogeneous part
- Find the annihilator  $a_2$  for the non-homogeneous part
- The annihilator for the whole recurrence is then  $a_1a_2$

13

## Another Example

- Consider  $T(n) = T(n-1) + T(n-2) + 2$ .
- The residue is  $\langle 2, 2, 2, \dots \rangle$  and
- The annihilator is still  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 1)$ , but the equation for  $T(2)$  changes!

14

## Another Example

- Consider  $T(n) = T(n-1) + T(n-2) + 2^n$ .
- The residue is  $\langle 1, 2, 4, 8, \dots \rangle$  and
- The annihilator is now  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 2)$ .

15

## Another Example

- Consider  $T(n) = T(n-1) + T(n-2) + n$ .
- The residue is  $\langle 1, 2, 3, 4, \dots \rangle$
- The annihilator is now  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 1)^2$ .

16

## Another Example

- Consider  $T(n) = T(n-1) + T(n-2) + n^2$ .
- The residue is  $\langle 1, 4, 9, 16, \dots \rangle$  and
- The annihilator is  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 1)^3$ .

17

## Another Example

- Consider  $T(n) = T(n-1) + T(n-2) + n^2 - 2^n$ .
- The residue is  $\langle 1 - 1, 4 - 4, 9 - 8, 16 - 16, \dots \rangle$  and the
- The annihilator is  $(\mathbf{L}^2 - \mathbf{L} - 1)(\mathbf{L} - 1)^3(\mathbf{L} - 2)$ .

18

## In Class Exercise

- Consider  $T(n) = 3 * T(n-1) + 3^n$
- Q1: What is the homogeneous part, and what annihilates it?
- Q2: What is the non-homogeneous part, and what annihilates it?
- Q3: What is the annihilator of  $T(n)$ , and what is the general form of the recurrence?

19

## Limitations

- Our method does not work on  $T(n) = T(n-1) + \frac{1}{n}$  or  $T(n) = T(n-1) + \lg n$
- The problem is that  $\frac{1}{n}$  and  $\lg n$  do not have annihilators.
- Our tool, as it stands, is limited.
- Key idea for strengthening it is *transformations*

20

## Transformations Idea

- Consider the recurrence giving the run time of mergesort  $T(n) = 2T(n/2) + kn$  (for some constant  $k$ ),  $T(1) = 1$
- How do we solve this?
- We have no technique for annihilating terms like  $T(n/2)$
- However, we can *transform* the recurrence into one with which we can work

21

## Transformation

- Let  $n = 2^i$  and rewrite  $T(n)$ :
- $T(2^0) = 1$  and  $T(2^i) = 2T(\frac{2^i}{2}) + k2^i = 2T(2^{i-1}) + k2^i$
- Now define a new sequence  $t$  as follows:  $t(i) = T(2^i)$
- Then  $t(0) = 1$ ,  $t(i) = 2t(i-1) + k2^i$

22

## Now Solve

- We've got a new recurrence:  $t(0) = 1$ ,  $t(i) = 2t(i-1) + k2^i$
- We can easily find the annihilator for this recurrence
- $(\mathbf{L} - 2)$  annihilates the homogeneous part,  $(\mathbf{L} - 2)$  annihilates the non-homogeneous part, So  $(\mathbf{L} - 2)(\mathbf{L} - 2)$  annihilates  $t(i)$
- Thus  $t(i) = (c_1i + c_2)2^i$

23

## Reverse Transformation

- We've got a solution for  $t(i)$  and we want to transform this into a solution for  $T(n)$
- Recall that  $t(i) = T(2^i)$  and  $2^i = n$

$$t(i) = (c_1 i + c_2) 2^i \quad (1)$$

$$T(2^i) = (c_1 i + c_2) 2^i \quad (2)$$

$$T(n) = (c_1 \lg n + c_2) n \quad (3)$$

$$= c_1 n \lg n + c_2 n \quad (4)$$

$$= \Theta(n \lg n) \quad (5)$$

24

## Success!

Let's recap what just happened:

- We could not find the annihilator of  $T(n)$  so:
- We did a *transformation* to a recurrence we could solve,  $t(i)$  (we let  $n = 2^i$  and  $t(i) = T(2^i)$ )
- We found the annihilator for  $t(i)$ , and solved the recurrence for  $t(i)$
- We *reverse transformed* the solution for  $t(i)$  back to a solution for  $T(n)$

25

## Another Example

- Consider the recurrence  $T(n) = 9T(\frac{n}{3}) + kn$ , where  $T(1) = 1$  and  $k$  is some constant
- Let  $n = 3^i$  and rewrite  $T(n)$ :
- $T(2^0) = 1$  and  $T(3^i) = 9T(3^{i-1}) + k3^i$
- Now define a sequence  $t$  as follows  $t(i) = T(3^i)$
- Then  $t(0) = 1$ ,  $t(i) = 9t(i-1) + k3^i$

26

## Now Solve

- $t(0) = 1$ ,  $t(i) = 9t(i-1) + k3^i$
- This is annihilated by  $(L-9)(L-3)$
- So  $t(i)$  is of the form  $t(i) = c_1 9^i + c_2 3^i$

27

## Reverse Transformation

- $t(i) = c_1 9^i + c_2 3^i$
- Recall:  $t(i) = T(3^i)$  and  $3^i = n$

$$\begin{aligned}t(i) &= c_1 9^i + c_2 3^i \\T(3^i) &= c_1 9^i + c_2 3^i \\T(n) &= c_1 (3^i)^2 + c_2 3^i \\&= c_1 n^2 + c_2 n \\&= \Theta(n^2)\end{aligned}$$

28

## In Class Exercise

Consider the recurrence  $T(n) = 2T(n/4) + kn$ , where  $T(1) = 1$ , and  $k$  is some constant

- Q1: What is the transformed recurrence  $t(i)$ ? How do we rewrite  $n$  and  $T(n)$  to get this sequence?
- Q2: What is the annihilator of  $t(i)$ ? What is the solution for the recurrence  $t(i)$ ?
- Q3: What is the solution for  $T(n)$ ? (i.e. do the reverse transformation)

29

## A Final Example

Not always obvious what sort of transformation to do:

- Consider  $T(n) = 2T(\sqrt{n}) + \log n$
- Let  $n = 2^i$  and rewrite  $T(n)$ :
- $T(2^i) = 2T(2^{i/2}) + i$
- Define  $t(i) = T(2^i)$ :
- $t(i) = 2t(i/2) + i$

30

## A Final Example

- This final recurrence is something we know how to solve!
- $t(i) = O(i \log i)$
- The reverse transform gives:

$$t(i) = O(i \log i) \quad (6)$$

$$T(2^i) = O(i \log i) \quad (7)$$

$$T(n) = O(\log n \log \log n) \quad (8)$$

31



## ┌ Todo ───

- HW 1
- Start Chapter 15 in text