

# CS 530: Geometric and Probabilistic Methods in Computer Science

## Homework 5 (Fall '11)

1. A bivariate Gaussian random variable,  $\mathbf{x} = \begin{bmatrix} x_0 & x_1 \end{bmatrix}^T$ , has the following p.d.f.:

$$f(x_0, x_1) = \frac{(a^2 - b^2)^{\frac{1}{2}}}{2\pi} \exp\left(-\frac{1}{2} [ax_0x_0 + 2bx_0x_1 + ax_1x_1]\right).$$

- (a) Give the matrix  $\mathbf{W}$  which will decorrelate the components of  $\mathbf{x}$ .
  - (b) Let  $\mathbf{u} = \mathbf{W}\mathbf{x}$ . Give an expression for  $g(u_0, u_1)$ , the p.d.f. for the bivariate Gaussian random variable,  $\mathbf{u} = \begin{bmatrix} u_0 & u_1 \end{bmatrix}^T$ .
2. Assume that the red, green, and blue components of the cactus image (cactus.ppm) found on the class homepage are samples of a  $3 \times 1$  vector random variable. Use MATLAB to do the following:
- (a) Compute the mean vector and subtract it from each of the pixels to produce a color image with zero mean.
  - (b) Compute the  $3 \times 3$  covariance matrix for the zero mean image.
  - (c) Compute the eigenvectors and eigenvalues of the covariance matrix.
  - (d) Transform each pixel in the cactus image using the matrix of eigenvectors of the covariance matrix (i.e., KL transform).
  - (e) Display the transformed image. If you have access to a color printer, provide hardcopy of the transformed image. Otherwise provide hardcopy of the red, green, and blue components of the transformed image.
  - (f) Show that the covariance matrix of the transformed image is diagonal. Show that the variances are equal to the eigenvalues of the covariance matrix of the untransformed image.

3. Prove the following:

$$\frac{\left[ \frac{P(x+\Delta x, t) - P(x, t)}{\Delta x} - \frac{P(x, t) - P(x - \Delta x, t)}{\Delta x} \right]}{\Delta x} = \frac{\partial^2 P}{\partial x^2} \Big|_{x, t} + O[(\Delta x^2)].$$

**Hint.** Replace  $P(x + \Delta x, t)$ , and  $P(x - \Delta x, t)$  with their Taylor series.

4. The partial differential equation (PDE) which governs the motion of waves propagating with velocity  $c$  in a two-dimensional medium is:

$$\frac{\partial^2 P}{\partial t^2} = c^2 \left[ \frac{\partial^2 P}{\partial x^2} + \frac{\partial^2 P}{\partial y^2} \right].$$

We will consider the evolution of functions of the torus. More precisely, we assume that  $P$  is periodic in  $x$  and  $y$  with periods,  $X$  and  $Y$ , i.e.,  $P(x, y) = P(x + jX, y + kY)$  for all integers,  $j$  and  $k$ .

- Using finite difference approximations for  $\partial^2 P / \partial t^2$ ,  $\partial^2 P / \partial x^2$ , and  $\partial^2 P / \partial y^2$ , derive an update equation for numerical solution of the two-dimensional wave equation, i.e., an equation for  $P_{x,y}^t$  in terms of  $P_{x,y}^{t-1}$ ,  $P_{x,y}^{t-2}$ ,  $P_{x-1,y}^{t-1}$ , etc.
- Implement and test your numerical scheme in MATLAB. Compute solutions of the wave equation for functions of a torus of size  $64 \text{ m} \times 64 \text{ m}$ . You should assume that  $\Delta x = \Delta y$  is  $1 \text{ m}$ ,  $\Delta t$  is  $0.005 \text{ s}$ , and the conduction velocity  $c$  is  $100 \text{ m/s}$ . The initial condition  $P_{x,y}^0 = P_{x,y}^1$  is a Gaussian of standard deviation,  $3 \text{ m}$ . For reasons of symmetry, the location of the mean is not important. Render your solutions as images (256 grey values) and provide hardcopy for  $t = 0.1 \text{ s}, 0.2 \text{ s}, \dots, 2.0 \text{ s}$ .
- **Hint.** The trickiest thing about this problem is properly computing the update on the boundaries of the periodic domain. This is relatively painless (and quite fast) if you recognize that the principal term in the update equation is actually a convolution which can be computed using the 2D FFT. For example, the second partial derivative in the  $x$  direction of a discrete  $N \times N$  image  $\mathbf{F}$  can be estimated at every location by convolving  $\mathbf{F}$  with the  $N \times N$  mask  $\mathbf{G}$ :

$$\left. \frac{\partial^2 F}{\partial x^2} \right|_{x,y} \approx \sum_{i=0}^{N-1} \sum_{j=0}^{N-1} F(i, j) \cdot G(x - i \bmod N, y - j \bmod N)$$

where

$$G = \begin{bmatrix} -2 & 1 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}.$$

Consequently,  $\frac{\partial^2 F}{\partial x^2} \approx \mathbf{F} * \mathbf{G} = \mathbf{W} (\hat{\mathbf{F}} \cdot \hat{\mathbf{G}}) \mathbf{W}$  where  $\hat{\mathbf{F}} = \mathbf{W}^* \mathbf{F} \mathbf{W}^*$  and  $\hat{\mathbf{G}} = \mathbf{W}^* \mathbf{G} \mathbf{W}^*$ .

If you choose not to use the FFT, you can still make your code fast by using vector operations instead of *for* loops whenever possible.

- **Optional stuff.** You can make your solutions colorful by rendering them as color images. One way to do this is to let  $P^t$  be red,  $P^{t-1}$  be green, and  $P^{t-2}$  be blue. You can animate your solutions using the Linux *animate* utility. Experiment with other values for  $c$ . At what velocity does the solution become numerically unstable? What does numerical instability look like?

5. A flying saucer containing an alien from Alpha Centauri has crashed in Roswell, NM. Analysis of the alien's visual system reveals it to be dichromatic with spectral sensitivity functions  $S_0(\lambda)$  and  $S_1(\lambda)$ . The wreckage of the spacecraft contains a flat panel display with two types of light emitting diodes. The spectral emission functions for these diodes are  $L_0(\lambda)$  and  $L_1(\lambda)$ . After interrogation of the alien by the Defense Threat Reduction Agency at their secret headquarters it has been learned that the alien is the vanguard of a species which is poised to invade Earth. The invasion can only be prevented by transmitting two voltages,  $v_0$  and  $v_1$ , which will reproduce the color of Alpha Centauri on the flat panel display of the flagship of the invasion fleet. Fortunately, the emission spectrum,  $A(\lambda)$ , of Alpha Centauri has been measured using Earth-based telescopes.

- Give the expressions for the cone photocurrents  $c_0(A)$  and  $c_1(A)$  evoked when an alien sees Alpha Centauri.
- Give the expression for the cone photocurrent  $c_0$  which is evoked from a weighted mixture of light sources with spectral emission functions,  $L_0$  and  $L_1$ , and weights,  $v_0$  and  $v_1$ .
- Give the expression for the color matching matrix,  $\mathbf{M}$ , which relates the cone photocurrents,  $c_0$  and  $c_1$ , to the voltages  $v_0$  and  $v_1$  required to reproduce the color on an alien flat panel display.
- Give expressions for  $v_0$  and  $v_1$ , the values which must be transmitted to prevent the destruction of Earth.