

## Marginal Probability Distribution

To compute  $p_X(k)$ , we sum  $p_{XY}(i, j)$  over pairs of  $i$  and  $j$  where  $i = k$ :

$$\begin{aligned} p_X(k) &= \sum_{\{i,j \mid i=k\}} p_{XY}(k, j) \\ &= \sum_{j=-\infty}^{\infty} p_{XY}(k, j). \end{aligned}$$

## Sum of Discrete r.v.'s

Let  $Z$  be a discrete random variable equal to the sum of the discrete random variables  $X$  and  $Y$ . To compute  $p_Z(k)$ , we sum  $p_{XY}(i, j)$  over pairs of  $i$  and  $j$  where  $i + j = k$ :

$$p_Z(k) = \sum_{\{i, j \mid i+j=k\}} p_{XY}(i, j).$$

Observe that  $i + j = k$  iff  $j = k - i$ . It follows that

$$p_Z(k) = \sum_{i=-\infty}^{\infty} p_{XY}(i, k - i)$$

If  $X$  and  $Y$  are statistically independent, then  $p_{XY}(i, j) = p_X(i)p_Y(j)$ . It follows that

$$p_Z(k) = \sum_{i=-\infty}^{\infty} p_X(i)p_Y(k - i).$$

## Sum of Discrete r.v.'s (contd.)

The *discrete convolution* of  $f$  and  $g$ , written  $f * g$ , is defined to be:

$$\{f * g\}(k) = \sum_{i=-\infty}^{\infty} f(i)g(k-i).$$

Accordingly, if  $Z = X + Y$ , then

$$p_Z = p_X * p_Y.$$

## Sum of Discrete r.v.'s (contd.)

Since  $i + j = k$  iff  $i = k - j$ , we could just as easily have written  $p_Z(k)$  as follows:

$$\begin{aligned} p_Z(k) &= \sum_{j=-\infty}^{\infty} p_{XY}(k-j, j) \\ &= \sum_{j=-\infty}^{\infty} p_X(k-j)p_Y(j). \end{aligned}$$

It follows that

$$p_X * p_Y = p_Y * p_X$$

and that convolution (like addition) is *commutative*.

## Difference of Discrete r.v.'s

Let  $Z$  be a discrete random variable equal to the difference of the discrete random variables  $X$  and  $Y$ . To compute  $p_Z(k)$ , we sum  $p_{XY}(i, j)$  over the pairs of  $i$  and  $j$  where  $i - j = k$ :

$$p_Z(k) = \sum_{\{i, j \mid i - j = k\}} p_{XY}(i, j).$$

Observe that  $i - j = k$  iff  $j = i - k$ . It follows that

$$p_Z(k) = \sum_{i=-\infty}^{\infty} p_{XY}(i, i - k).$$

If  $X$  and  $Y$  are statistically independent, then  $p_{XY}(i, j) = p_X(i)p_Y(j)$ . It follows that

$$p_Z(k) = \sum_{i=-\infty}^{\infty} p_X(i)p_Y(i - k).$$

## Difference of Discrete r.v.'s (contd).

The *discrete correlation* of  $f$  and  $g$ , is defined to be:

$$\{f * g(-(\cdot))\}(k) = \sum_{i=-\infty}^{\infty} f(i)g(i-k).$$

## Marginal Probability Density

To compute  $f_X(z)$ , we integrate  $f_{XY}(x, y)$  along the line  $x = z$ :

$$f_X(z) = \int_{-\infty}^{\infty} f_{XY}(z, y) dy$$

## Sum of Continuous r.v.'s

Let  $Z$  be a continuous random variable equal to the sum of the continuous random variables,  $X$  and  $Y$ . To compute  $f_Z(z)$ , we integrate  $f_{XY}(x, y)$  along the line  $x + y = z$  or  $y = z - x$ :

$$f_Z(z) = \int_{-\infty}^{\infty} f_{XY}(x, z - x) dx.$$

If  $X$  and  $Y$  are statistically independent, then  $f_{XY}(x, y) = f_X(x)f_Y(y)$ . It follows that

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z - x) dx.$$

## Sum of Continuous r.v.'s (contd.)

The *convolution* of  $f$  and  $g$ , written  $f * g$ , is defined to be

$$\{f * g\}(v) = \int_{-\infty}^{\infty} f(u)g(v - u)du.$$

Accordingly, if  $Z = X + Y$ , then

$$f_Z = f_X * f_Y = f_Y * f_X.$$

## Difference of Continuous r.v.'s

Let  $Z$  be a continuous random variable equal to the difference of the continuous random variables,  $X$  and  $Y$ . To compute  $f_Z(z)$ , we integrate  $f_{XY}(x, y)$  along the line where  $x - y = z$  or  $y = x - z$ :

$$f_Z(z) = \int_{-\infty}^{\infty} f_{XY}(x, x - z) dx.$$

If  $X$  and  $Y$  are statistically independent, then  $f_{XY}(x, y) = f_X(x)f_Y(y)$ . It follows that

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(x - z) dx.$$

## Difference of Continuous r.v.'s (contd.)

The *correlation* of  $f$  and  $g$ , is defined to be

$$\{f * g(-(\cdot))\}(v) = \int_{-\infty}^{\infty} f(u)g(u - v)du$$

Accordingly, if  $Z = X - Y$ , then

$$f_Z(z) = \{f_X * f_Y(-(\cdot))\}(z)$$

## Law of Large Numbers

Let  $X_1, X_2, \dots, X_N$  be samples of a random variable,  $X$ . It follows that  $X_1, X_2, \dots, X_N$  are independent, identically distributed (i.i.d.) random variables. Then

$$\lim_{N \rightarrow \infty} \frac{X_1 + X_2 + \dots + X_N}{N} = \mu = \langle X \rangle.$$

i.e., the mean of an infinite number of samples of a random variable equals the expected value.

## Central Limit Theorem

Let  $X_1, X_2, \dots, X_N$  be samples of a random variable,  $X$ . It follows that  $X_1, X_2, \dots, X_N$  are independent, identically distributed (i.i.d.) random variables. Then

$$\lim_{N \rightarrow \infty} P \left( \frac{X_1 + X_2 + \dots + X_N - N\mu}{\sigma\sqrt{N}} \leq a \right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx.$$

i.e., the sum of an infinite number of i.i.d. random variables is a random variable with Gaussian density.

## Central Limit Theorem (contd.)

Multiplying numerator and denominator by  $\frac{1}{N}$  yields:

$$\lim_{N \rightarrow \infty} P \left( \frac{\frac{X_1 + X_2 + \dots + X_N}{N} - \mu}{\sigma / \sqrt{N}} \leq a \right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx.$$