

CS 506, HW 2

Prof. Jared Saia, University of New Mexico

You are encouraged to work on the homework in groups of about 2 or 3. You may turn in one writeup per group, but please include a paragraph stating specifically in which way group members contributed for each problem.

1. Let P be a set of n points in \mathbb{R}^2 . Show that P can be preprocessed in $O(n^2 \log n)$ time to create a data structure that can answer any query of the type: “How many points lie below this query line?” in $O(\log n)$ time.
2. The following question is about maximum degree vertices in Voronoi diagrams and triangulations.
 - (a) Prove that for any $n > 3$ there is a set of n points in the plane such that one of the cells of the Voronoi diagram of these points has $n - 1$ vertices.
 - (b) The degree of a point in a triangulation is the number of edges incident to it. Give an example of a set of n points in the plane, such that, no matter *how* the set is triangulated, there is always a point whose degree is $n - 1$.
3. Prove Euler’s formula for the Euclidean plane. In particular, prove that for any connected planar graph, $V - E + F = 2$ where V is the number of vertices, E is the number of edges, and F is the number of faces in the planar graph. Hint: Use proof by induction. There are multiple quantities that you can do induction on; the fun part of this problem is to find the right quantity.
4. Recall that a 5-clique is a graph of 5 nodes that are all completely connected
 - (a) Prove that you can not draw a 5-clique in the plane with no edge crossings. Hint: Use Euler’s formula.

- (b) The Euler characteristic of a torus is 0 (recall this means that $V - E + F = 0$, for any graph drawn on a torus, where V is vertices, E is edges and F is faces). Can you draw a 5-clique with no edge crossings in (1) the Euclidean plane; and (2) a torus? In both cases, if the answer is yes, please demonstrate. If the answer is no, prove it is impossible by using Euler's formula, counting the values that would be needed for V , E and F , and showing a contradiction.¹
- (c) The graph $K_{3,3}$ is the complete bipartite graph with 3 nodes on both sides. It's the 3 home, 3 utility graph that we proved can't be drawn with no edge crossings in the plane. A Mobius strip² has Euler characteristic 0. Can you draw $K_{3,3}$ with no edge crossings on a Moibus strip? If so, please demonstrate (google Moibus strip for how to make one with tape and scissors). If not, prove it is impossible using Eurler's identity.
5. (Based on Exercise 9.16 from Berg) A k -clustering of a set P of n points in the the plane is a partitioning of P into k non-empty subsets P_1, \dots, P_k . Define the distance between any pair P_i, P_j of clusters to be the minimum distance between one point from P_i and one point from P_j . In particular,;

$$dist(P_i, P_j) = \min_{p \in P_i, q \in P_j} dist(p, q)$$

We want to find a k -clustering (for given k and P) that maximizes the minimum distance between clusters.

- (a) Suppose the minimum distance between any pair of clusters is achieved by points $p \in P_i$ and $q \in P_j$, for some clusters P_i and P_j . Prove that edge \overline{pq} is an edge of the Delaunay triangulation of P
- (b) Give an $O(n \log n)$ time algorithm to compute a k -clustering maximizing the minimum distance between clusters. Hint: Use a Union-Find data structure!

¹For examples of how to show a drawing for a torus and a Mobius strip, see: https://ptwiddle.github.io/Graph-Theory-Notes/s_graphs_on_surfaces_other_surfaces.html

²Fun fact: Mobius strips are sometimes used in conveyer belts to ensure the entire surface area of the belt gets even wear.

6. Write a brief proposal for your class project (1 page). Please talk to me about your ideas briefly before you start writing this.